

Acute effects of endocannabinoid anandamide and CB1 receptor antagonist, AM251 in the regulation of thyrotropin secretion

Marco Aurélio Liberato Costa da Veiga¹, Flávia Fonseca Bloise¹, Ricardo Henrique Costa-e-Sousa¹, Luana Lopes Souza¹, Norma Aparecida dos Santos Almeida¹, Karen Jesus Oliveira^{1,2} and Carmen Cabanelas Pazos-Moura¹

¹Laboratório de Endocrinologia Molecular, Instituto de Biofísica Carlos Chagas Filho, Universidade Federal do Rio de Janeiro, Rio de Janeiro 21949-900, Brazil

²Departamento de Fisiologia e Farmacologia, Universidade Federal Fluminense, Rio de Janeiro 23070-200, Brazil

(Correspondence should be addressed to C C Pazos-Moura; Email: cpazosm@biof.ufrj.br)

Abstract

We examined the acute effects of endocannabinoid, anandamide, and of synthetic cannabinoid receptor antagonist, AM251 [*N*-(piperidin-1-yl)-1-(2,4-dichlorophenyl)-5-(4-chlorophenyl)-4-methyl-1*H*-pyrazole-3-carboxamide], on TSH, thyroxine (T₄), and triiodothyronine (T₃) secretions. Euthyroid male rats showed a 42% decrease in serum TSH, 2 h after a single i.p. injection of 0.02, but not 0.2 mg/kg body weight (BW), anandamide, accompanied by a 39% reduction in serum T₄, without alteration in serum T₃. At 0.5 and 1 h, these serum hormones showed no significant change. Hypothyroid rats showed a 35% reduction in serum TSH ($P < 0.01$), 2 h after anandamide injection, which had no effect on hyperthyroid rats. In both thyroid states, no modification of serum thyroid hormones was observed. Intraperitoneal injection of 0.17 or 1.7 mg/kg BW AM251 in euthyroid rats caused, 1.5 h later, 1.7-fold or 4.3-fold increase in serum TSH respectively, without changing thyroid

hormones. Stimulatory effect of 0.17 mg/kg BW AM251 and inhibitory effect of anandamide was abolished in the group injected with AM251 followed by an anandamide injection, 30 min later. Intracerebroventricular injection of 20 ng (but not 200 ng) anandamide induced a decrease in serum TSH at 60 min after injection, which tended to disappear at 120 min. Anterior pituitary explants presented significant reduction in TSH release in the presence of 10^{-7} M anandamide in incubation medium, which was blocked by 10^{-7} M AM251. In conclusion, anandamide has the ability to acutely inhibit TSH release in eu- and hypothyroid rats, acting at the hypothalamus–pituitary axis. Since, in addition, the cannabinoid receptor antagonist AM251 increased TSH release, we suggest that endocannabinoid system has a role as negative regulator of TSH secretion.

Journal of Endocrinology (2008) **199**, 235–242

Introduction

The active principle isolated from *Cannabis sativa*, Δ^9 -tetrahydrocannabinol (Δ^9 -THC), has many well-known pharmacological properties. The actions of this phytocannabinoid are mediated by two subtypes of G-protein coupled receptors, CB1 (also known as CNR1) and CB2 (also known as CNR2) (Devane *et al.* 1988, Matsuda *et al.* 1990, Munro *et al.* 1993). CB1 is abundant in the central nervous system, and it is also present in the peripheral nervous system and in several peripheral organs (Tsou *et al.* 1998). By contrast, CB2 receptor expression in the brain is very low and it is located mostly in immune cells (Klein 2005). Endogenous ligands of cannabinoid receptors were identified and named endocannabinoids (Di Marzo & Fontana 1995). The main endocannabinoids are arachidonoyl ethanolamide (anandamide) and 2-arachidonoyl glycerol, derivatives of arachidonic acid, which are produced by the cleavage of membrane lipid precursors (Di Marzo *et al.* 2004). Endocannabinoids and

their receptors together with enzymes for ligand biosynthesis and inactivation comprise the endocannabinoid system (Di Marzo *et al.* 2004). This system participates in several physiological processes, such as nociceptive transmission, motor activity, learning and memory process, emotional responses, as well as thermoregulation and appetite regulation (Di Marzo *et al.* 2004, Foster-Schubert & Cummings 2006).

In addition, it has been shown that endocannabinoids have a role in the control of the neuroendocrine system. Anandamide modulates prolactin (PRL) and luteinizing hormone (LH) CGR secretions in a sex steroid-dependent manner. In male rats and female ovariectomized rats, anandamide had the ability to suppress PRL and LH release (Wenger *et al.* 1995, Scorticati *et al.* 2003, 2004), while in ovariectomized estrogen-primed rats, it stimulated PRL and LH release (Scorticati *et al.* 2003, 2004). Regarding growth hormone secretion, one report found that Δ^9 -THC administration increased while anandamide decreased the hormone serum concentrations (Wenger *et al.* 1995). There are also reports that anandamide,

centrally injected in rats, was able to increase the secretion of adrenocorticotropin hormone (Weidenfeld *et al.* 1994).

Earlier reports have demonstrated the ability of a single injection of the phytocannabinoid Δ^9 -THC to reduce serum concentrations of TSH and thyroid hormones (Hillard *et al.* 1984). Another study (Lomax *et al.* 1970) suggested that the reduction in thyrotropic axis would be consequent to hypothalamic action of Δ^9 -THC, resulting in decreased thyrotropin-releasing hormone (TRH) release, although so far, direct evidence is lacking. More recently, it was shown that the synthetic cannabinoid receptor agonist WIN 55212-2 also reduced serum triiodothyronine (T_3) and thyroxine (T_4), but no alteration was observed on serum TSH (Porcella *et al.* 2002). Therefore, in the present paper, in order to contribute further insights into the role of the endogenous cannabinoid system on thyrotropic axis function, we investigated the acute effect of anandamide on TSH secretion *in vivo* and *in vitro*. We also analyzed the acute effect of a CB1 receptor antagonist, AM251, since CB1 receptors are present at hypothalamus, pituitary, and thyroid glands (Fernandez-Ruiz *et al.* 1997, Wenger *et al.* 1999, Porcella *et al.* 2002, Scorticati *et al.* 2004).

Materials and Methods

Animals

Adult male Wistar rats, 250–300 g body weight (BW), were kept under a 12 h light:12 h darkness cycle (light onset at 0700 h) at 24 ± 1 °C; a standard diet and tap water were available *ad libitum*. All experimental protocols were approved by our institutional animal care committee Comissão de Avaliação de Utilização de Animais em Pesquisa (CAUAP) and are in accordance with the FRAME's guidelines. Experiments were performed with eu-, hypo-, and hyperthyroid rats. Hypothyroidism was induced by treatment with 0.03% methimazole (Sigma) in the drinking water for 21 days. Hyperthyroidism was induced by s.c. daily single injection of T_4 (L-Thyroxine Sigma), 10 μ g/100 g BW for 6 days.

In vivo experiments

Systemic administration of anandamide or AM251, a selective CB1 receptor antagonist Euthyroid animals were divided into four groups that received a single i.p. injection of vehicle (propylene glycol:ethanol:saline 1:1:2 – control) or 0.02 mg/kg BW anandamide (Tocris Cookson Inc., Ellisville, MO, USA) and killed by decapitation 0.5, 1, or 2 h after the injection. Control group was killed at a time between 1 and 2 h. Other group of euthyroid rats received a single i.p. administration of vehicle or 0.2 mg/kg BW anandamide and killed 2 h later. Drug administration protocol was based on that employed by Wenger *et al.* (1997) who demonstrated that 0.02 mg/kg BW anandamide stimulated the secretion of adrenocorticotropin hormone (ACTH) and corticosterone in rats.

Hypo- and hyperthyroid animals were divided into two groups that received a single i.p. injection of vehicle (control) or 0.02 mg/kg BW anandamide, and killed 2 h later. The time was chosen based on the results observed on euthyroid animals.

In order to evaluate the response of the thyroid axis to selective CB1 receptor antagonist/inverse agonist, AM251, euthyroid animals were injected i.p. with vehicle (3% DMSO), 0.17 mg/kg or 1.7 mg/kg BW of AM251 [*N*-(piperidin-1-yl)-1-(2,4-dichlorophenyl)-5-(4-chlorophenyl)-4-methyl-1*H*-pyrazole-3-carboxamide] (Tocris Cookson Inc.), and killed 1.5 h later. AM251 dose was chosen based on previous reports on behavior studies that had injected rats with the drug intraperitoneally (Chambers *et al.* 2004, McLaughlin *et al.* 2005, Xi *et al.* 2006).

In order to investigate the involvement of endocannabinoid receptors on the effect of anandamide on the thyrotropic axis, 0.17 mg/kg BW AM251 was injected 30 min before the administration of 0.02 mg/kg anandamide in euthyroid rats, and compared with groups treated with 0.02 mg/kg anandamide alone or 0.17 mg/kg BW AM251 alone. To serve as a control group, half of the animals were injected with AM251 vehicle and other half with anandamide vehicle, and since data of these two groups did not differ, they were pooled and presented together. Animals were killed 2 h after anandamide injection.

In all experiments, animals were killed by decapitation and serum was obtained from trunk blood to measure TSH and total T_4 and T_3 concentrations. Experiments were performed until 1100 h, in order to avoid circadian fluctuations interfering with results.

Central administration of anandamide

Rats underwent third ventricular cannulation (i.c.v.) 7 days before the experiment. A stainless steel guide cannula (0.6 mm o.d.) for microinjection was implanted 0.5 mm posterior to bregma and 8.0–8.5 mm ventral to the skull, as reported before (Antunes-Rodrigues & McCann 1970, Rettori *et al.* 1992), under ketamine plus xylazine anesthesia. Verification of correct cannula placement was made by the observation of fluid-filled column (attached to guide cannula) displacement. The day before the experiment, a jugular catheter was implanted, under the same anesthetic protocol described above. Briefly, the jugular vein was isolated by blunt dissection and a catheter made of silastic tubing was introduced about 2.2 mm down. The free end of the catheter was passed under the skin and externalized between the scapulae. The catheter was flushed with heparin saline (50 U/ml). The next morning, extension tubing was attached to the jugular catheter and animals were left undisturbed for half an hour, then a basal heparinized blood sample was collected, followed by the i.c.v. injection of a solution of 20 or 200 ng anandamide in 2 μ l or an equal volume of vehicle (Trocrisolve, Tocris Cookson Inc). Blood samples were collected 30, 60, and 120 min after microinjection. After each sample collection, volume replacement was conducted with sterile saline.

In vitro experiments: pituitary explants

The *in vitro* protocol was performed as described previously (Rettori *et al.* 1989, 1992, Pazos-Moura *et al.* 1996, Ortiga-Carvalho *et al.* 2002). Euthyroid animals were killed by decapitation and their anterior pituitaries were quickly dissected. Each hemi-pituitary was immediately transferred to a flask containing 1 ml Krebs–Ringer bicarbonate medium (pH 7.4) at 37 °C in an atmosphere of 95% O₂:5% CO₂ in a Dubnoff metabolic shaker. After a 30-min preincubation period, medium was changed and glands were incubated in the absence or presence of tested drugs. In the first experiment, glands were incubated with 10⁻¹¹, 10⁻⁹, or 10⁻⁷ M anandamide in Tocrisolve (Tocris Cookson Inc). In the second experiment, glands were incubated with anandamide or AM251 or both drugs at the same molarity (10⁻⁷ M). At the end of 2-h incubation an aliquot was removed, and kept at -20 °C until assayed for TSH measurement.

Hormone measurements

TSH concentrations in serum and in incubation medium were measured by specific RIA, employing reagents supplied by National Hormone and Peptide Program (Torrance, CA, USA), as previously described (Chard 1987, Ortiga-Carvalho *et al.* 1996), and it was expressed in terms of the reference preparation (RP3). The assay is highly specific for rat TSH and no significant cross-reactivity is reported by the supplier. Minimum assay detection value was 0.18 ng/ml and within-assay variation was less than 8% in all assays. Total serum T₄ and T₃ was detected by RIA (Total T₄ or Total T₃ Mab-ICN Pharmaceuticals, Costa Mesa, CA, USA). Minimum assay detection value was 25 ng/dl for T₃, 1 µg/dl for T₄, and within-assay variation was less than 4.5% for T₃ and less than 3.5% for T₄ in all assays. Samples of the same experiment were run within the same assay.

Statistical analyses

Data are reported as mean ± s.e.m. Serum TSH was analyzed employing Kruskal–Wallis test followed by Dunn's multiple comparison test or Mann–Whitney test, in experiments with two experimental groups. One-way ANOVA followed by Dunnett's multiple comparison test was employed for an assessment of the significance of *in vitro* TSH. T₄ and T₃ data were analyzed by ANOVA or unpaired *t*-test, according to the number of experimental groups (Zar 1996). Null hypothesis was discarded when *P* < 0.05.

Results

Acute effect of anandamide on thyrotropin and thyroid hormones secretion on euthyroid, hypothyroid, and hyperthyroid rats

In euthyroid rats (Fig. 1A), serum TSH was decreased by ~42% in animals that had received, 2 h earlier, a single




Figure 1 Serum (A) TSH, (B) T₄ and T₃ concentrations in euthyroid rats after i.p. administration of 0.02 mg/kg BW anandamide (ANA) or vehicle (C, control). Determinations were performed at 0.5, 1, and 2 h after the injection. *n* = 10–12 animals per group. **P* < 0.05 ***P* < 0.01 versus control group. Data are presented as mean ± s.e.m.

systemic injection of 0.02 mg/kg BW anandamide (*P* < 0.05). At the same time interval, serum T₄ showed a 39% reduction (*P* < 0.01; Fig. 1B). At earlier time-points (0.5 and 1 h), anandamide had no significant effect on serum TSH or serum T₄. Serum T₃ was statistically similar among all groups (Fig. 1B). However, a ten-time higher anandamide dose (0.2 mg/kg) had no effect on serum TSH, T₄, and T₃ (Fig. 2).

In hypothyroid animals, 2 h after the low-dose-anandamide administration, a reduction of ~35% in serum TSH was




Figure 2 Serum (A) TSH, (B) T₄ and T₃ concentrations in euthyroid rats after i.p. administration of 0.2 mg/kg BW anandamide (ANA 0.2) or vehicle (C, control). Determinations were performed at 2 h after the injection. *n* = 9–10 animals per group. Data are presented as mean ± s.e.m.

observed ($P < 0.01$; Fig. 3A), but there was no change in serum thyroid hormones concentrations (Fig. 3B).

In hyperthyroid animals, serum TSH did not change in response to low dose anandamide administration. As expected, hyperthyroid rats presented reduced serum TSH (0.4 ± 0.27 ng/ml for vehicle; $n = 4$ and 0.3 ± 0.1 ng/ml for anandamide treated; $n = 3$) and some of the animals (five of vehicle- and four of anandamide-treated groups) had serum TSH lower than the detection limit of assay (0.18 ng/ml). Also, serum T_4 and T_3 remained unchanged by anandamide treatment (T_4 : 7.85 ± 0.4 μ g/dl for vehicle, and 8.08 ± 0.4 μ g/dl for anandamide injected; T_3 : 132.9 ± 9.1 for vehicle and 154.8 ± 9.9 ng/dl for anandamide treatment).

Acute effect of CB1 receptor antagonist AM251 on thyrotropin secretion of euthyroid rats

As depicted in Fig. 4A, a single injection of AM251, injected 1.5 h before killing, induced a dose increase in serum TSH, of 1.7 ($P < 0.05$) or 4.3 times ($P < 0.001$) in animals receiving 0.17 or 1.7 mg/kg BW AM251 respectively (Fig. 4B). However, no statistically significant effects were observed regarding serum T_4 and T_3 .

Acute effect of CB1 receptor antagonist AM251 and anandamide on thyrotropin secretion of euthyroid rats

Figure 5 shows that the inhibitory effect of 0.02 mg/kg BW anandamide and the stimulatory effect of 0.17 mg/kg BW AM251 were abolished in rats receiving AM251 30 min before the injection of anandamide. The latter group presented serum TSH statistically similar to control group. All animals were killed 2 h after anandamide injection.

Effect of intracerebroventricular injection of anandamide in euthyroid rats

Serum TSH was evaluated before and at 30, 60, and 120 min after injection into the third ventricle of 20 or 200 ng anandamide. Figure 6 shows the variation of serum TSH in relation to basal levels (before injection). At 30 min,




Figure 3 Serum (A) TSH, (B) T_4 and T_3 concentrations in hypothyroid rats after i.p. administration of 0.02 mg/kg BW anandamide (ANA 0.02) or vehicle (C, control). Determinations were performed at 2 h after the injection. $n = 8-9$ animals per group. * $P < 0.01$ versus control group. Data are presented as mean \pm S.E.M.




Figure 4 Serum (A) TSH, (B) T_4 and T_3 concentrations in euthyroid rats after i.p. administration of the selective CB1 receptor antagonist, AM251 in the dose of 0.17 or 1.7 mg/kg BW or vehicle (C, control). Determinations were performed at 1.5 h after the injection. $n = 10-11$ animals per group. $P < 0.05$ versus *control group or versus #AM251 0.17. Data are presented as mean \pm S.E.M.

anandamide injection, at both doses, caused no significant change in TSH secretion. At 60 min, anandamide at 20 ng induced a significant reduction in TSH secretion ($P < 0.05$). The same trend was observed with 200 ng anandamide; however, the variation in serum TSH was not statistically significant. One hour later (at 120 min after microinjection), no significant effect was observed, although only the group treated with 20 ng anandamide still had a trend to reduce TSH secretion.

Effect of anandamide and AM251 in pituitary explants of euthyroid rats

The presence of anandamide in incubation media of hemipituitary glands showed a dose-related effect on TSH released to the medium. At 10^{-11} M, release of TSH was increased (1.6 times, $P < 0.05$); however, at 10^{-7} M, there was a significant reduction (66%), while the intermediary dose (10^{-9} M) induced no significant change in secretion




Figure 5 Serum (A) TSH, (B) T₄ and T₃ concentrations in euthyroid rats after i.p. administration of 0.02 mg/kg BW anandamide (ANA 0.02) or 0.17 mg/kg BW AM251, a selective CB1 receptor antagonist (AM251 0.17), or 0.17 mg/kg BW AM251 followed by 0.02 mg/kg BW anandamide injection 30 min later (ANA +AM251) or vehicle (C, control). Determinations were performed at 2 h after ANA injection. *n* = 9–11 animals per group. *P* < 0.05 versus *control group versus #ANA 0.02. Data are presented as mean ± S.E.M.

(Fig. 7A). In the second experiment, we observed that the inhibitory effect of 10⁻⁷ M anandamide in TSH release was blocked by the CB1 receptor antagonist AM251 at 10⁻⁷ M (*P* < 0.05; Fig. 7B). The incubation with 10⁻⁷ M AM251 alone showed a non-statistically significant trend to increase TSH release in comparison with control and a significant increase in relation to anandamide-treated group (*P* < 0.05), which showed a 30% decrease in relation to control (*P* < 0.05; Fig. 7B).

Discussion

The present study demonstrated that the endocannabinoid, anandamide, injected systemically or centrally into rats exhibited the ability to inhibit TSH secretion, while acute administration of the inverse agonist AM251 increases the hormone secretion.

The inhibitory effect of anandamide on TSH secretion occurred 2 h after systemic injection and was accompanied by concomitant reduction in serum T₄. These data suggest that anandamide acts at the pituitary and/or hypothalamus in order to inhibit TSH secretion, since increased TSH would be expected if the only site of action was the thyroid. An earlier study showed that the phytocannabinoid Δ9-THC was able to reduce serum TSH from 15 min to 3 h after a single i.p. injection, followed by a decrease in serum T₃ and T₄ at 3 and 6 h respectively (Hillard *et al.* 1984). However, in our study, the reduction in serum TSH and T₄ was concomitant and, considering the long half-life of serum T₄, unless TSH bioactivity was reduced at a much earlier time, it is likely that anandamide was also acting at the thyroid gland. In support to a direct action in the thyroid, there was the previous report by Porcella *et al.* (2002), showing that 4 h after the injection of cannabinoid receptor agonist WIN 55212-2, serum thyroid hormones were reduced without changes in serum TSH. This effect of the




Figure 6 Serum TSH variation (ΔTSH) in relation to zero time, before microinjection, and at 30, 60, or 120 min after microinjection of 20 or 200 ng anandamide into the third cerebral ventricle of euthyroid rats. Data represent mean ± S.E.M., *n* = 6–8 animals per group. **P* < 0.05 versus control group.




Figure 7 (A) TSH release from isolated rat hemi-anterior pituitary glands incubated in the absence (C, control) or presence of anandamide at 10^{-11} , 10^{-9} , or 10^{-7} M for 2 h. (B) TSH release from isolated rat hemi-anterior pituitary glands incubated in the absence (C, control), presence of 10^{-7} M anandamide (ANA 10^{-7}), 10^{-7} M AM251 (AM251 10^{-7}), or both drugs at the same molarity (ANA+AM251) for 2 h. Data represent mean \pm S.E.M. and correspond to two experiments and expressed in relation to basal secretion, which was set to 1. $n=9-14$, hemi-pituitaries/group. $P<0.05$ versus *all groups and versus #ANA 10^{-7} .

agonist was abolished by pre-treatment with specific CB1 antagonist SR 141716A. However, it is not clear which subtype of cannabinoid receptor mediates the effect of WIN 55212-2, on thyroid, since, so far, only CB1 was found in the thyroid (Porcella *et al.* 2002), and WIN 55212-2 is a CB2 preferring agonist.

Further evidence of the central action of anandamide was obtained by the demonstration of the inhibitory anandamide effect of on TSH secretion, without changes in serum thyroid hormones, in rats treated with methimazol, a drug that inhibits thyroid hormone biosynthesis. In addition, the fact that anandamide preserves its ability to reduce serum TSH in hypothyroid state highlights the potency of endocannabinoid effect on TSH regulation, since lack of thyroid hormones is a major determinant of high serum TSH (Weintraub *et al.*

2001). However, apparently, anandamide was not able to further decrease the already suppressed TSH serum concentrations of hyperthyroid rats.

The involvement of CB1 receptors in the control of TSH release is suggested by the acute rise in serum TSH induced by AM251 and by the fact that AM251 pre-treatment prevented anandamide effect. AM251 stimulatory effect on TSH release may be interpreted as that of anandamide, which acts preferentially by CB1 receptors, having a tonic inhibitory effect that is blocked by the CB1 receptor antagonist. Another likely explanation would be that AM251, acting as an inverse agonist, reduces the CB1 receptor signaling transduction pathway (Pertwee 2005). AM251, injected into the third cerebral ventricle, had been reported to change LH and PRL secretions in the opposite direction to anandamide effects (Scorticati *et al.* 2003, 2004). Serum thyroid hormones did not change in response to the rise in TSH induced by AM251, which may be related to the short time observed after the drug administration (1.5 h) or to the biological potency of the amount of TSH secreted. In a previous study (Porcella *et al.* 2002), systemic injection of CB1 receptor antagonist SR141716A did not modify serum TSH, T_3 , or T_4 . However, hormonal evaluation was performed 4 h after injection, while we used an earlier time in the current study. Therefore, it remains to be elucidated whether these different reports are related to time-course or to particularities attributed to each drug.

CB1 receptors have been demonstrated, albeit in low density, in hypothalamic nuclei (Fernandez-Ruiz *et al.* 1997, Scorticati *et al.* 2004) and also CB1-like immunoreactivity was found in rat and human anterior pituitaries (Fernandez-Ruiz *et al.* 1997, Pagotto *et al.* 2001, Scorticati *et al.* 2004). Therefore, the *in vivo* effect of anandamide on TSH release may result from a direct action at pituitary and/or hypothalamus. The possibility of a direct hypothalamic action is reinforced by the inhibitory effect we observed within 60 min after the i.c.v. administration of 20 ng anandamide. Wenger *et al.* (1997) had used a systemic treatment regimen identical to that employed in the present study and demonstrated that anandamide increased POMC release in concomitance with proto-oncogene protein c-fos (FOS) activation in paraventricular nucleus (PVN) hypothalamic neurons, peaking at 90 min after anandamide injection. Therefore, the time-course of our results is in accordance with a postulated action at hypothalamic site. In addition, earlier studies had shown that the phytocannabinoid Δ^9 -THC was not able to acutely affect serum TSH in animals with electrolytic lesions in medial mammillary nuclei of the hypothalamus (Lomax *et al.* 1970). Regarding systemic administration of AM251, a hypothalamic site of action is also possible, since previous studies have shown that following i.v. injection of [123 I]AM251 in mice, the peak whole-brain uptake of about 1% of the administered activity occurred at about 1 h and declined thereafter (Gatley *et al.* 1996). In addition, there are reports of centrally mediated effects induced by systemic acute administration of AM251,

employed in the same dose range of our higher dose, such as a decrease in food ingestion behavior (Chambers *et al.* 2004) and inhibition of cocaine-induced reinstatement of reward-seeking behavior (Xi *et al.* 2006).

The precise mechanism by which endogenous cannabinoid may modify TSH release cannot be elucidated by the present study. Considering the short half-life of anandamide and the relative long latency for the anandamide central effect on serum TSH, it is possible that this endocannabinoid did not act directly on TRH neurons, but likely that other mechanisms are involved (de Miguel *et al.* 1998, Scorticati *et al.* 2003). Scorticati *et al.* (2003) reported that dopamine concentration in medial basal hypothalamus and anterior pituitary was increased 90 min after a single intracerebroventricular injection of anandamide, which concomitantly induced inhibition of PRL release (Scorticati *et al.* 2003). Since dopamine is an inhibitor of PRL and TSH (Weintraub *et al.* 2001), a mechanism by which anandamide may inhibit PRL and also TSH release is via increasing dopamine input.

However, endocannabinoids may also modulate TSH secretion by a direct pituitary action. We were able to demonstrate the ability of anandamide at 10^{-7} M to reduce TSH release, an effect at least in part mediated by cannabinoid receptor 1, since it was blocked by AM251. However, anandamide at lower molarity (10^{-11} M) provoked an increase in TSH secretion to medium, and therefore the pituitary action of endogenous cannabinoid may vary depending on the concentration present in the gland.

Interestingly, a ten-time higher dose of anandamide, injected systemically or centrally, was not able to affect the thyrotropic axis. There are many reports showing lower biological efficacy of endogenous substances administered at higher dose than lower doses, including anandamide (Ortiga-Carvalho *et al.* 2002, Fernandez-Solari *et al.* 2004, Scorticati *et al.* 2004, Oliveira *et al.* 2006).

Thyroid hormones and endocannabinoids participate in energy homeostasis in opposite directions; while thyroid hormones are major stimulants of the basal metabolic rate and energy expenditure, endocannabinoids are important stimulators of food ingestion and energy conservation. During starvation, thyrotropic axis activity is reduced (Weintraub *et al.* 2001), while endocannabinoid levels increase (Matias & Di Marzo 2007). Therefore, it seems feasible to the existence of a physiologically relevant negative regulation of thyrotropic axis by endocannabinoids. In addition, since CB1 receptor antagonists are currently being introduced in therapeutics (Di Marzo *et al.* 2004), it will be important to investigate the consequences to pituitary–thyroid axis of chronic administration of CB1-receptors antagonists.

In conclusion, anandamide, an endogenous cannabinoid, has the ability to acutely inhibit TSH release in eu- and hypothyroid rats, acting at the hypothalamus–pituitary axis. Since the cannabinoid receptor antagonist, AM251, increased TSH release *in vivo* and blocked the effect of anandamide, it is suggested that endocannabinoid system may play a role as a negative regulator of TSH secretion *in vivo*.

Declaration of interest

The authors declare that there is no conflict of interest that would prejudice the impartiality of this scientific work.

Funding

This work was supported by Conselho Nacional de Pesquisa e Desenvolvimento and Fundação Carlos Chagas Filho de Amparo a Pesquisa do Estado do Rio de Janeiro.

References

- Antunes-Rodrigues J & McCann SM 1970 Water, sodium chloride, and food intake induced by injections of cholinergic and adrenergic drugs into third ventricle of the rat brain. *Proceedings of the Society for Experimental Biology and Medicine* **133** 1464–1470.
- Chambers AP, Sharkey KA & Koopmans HS 2004 Cannabinoid (CB1) receptor antagonist, AM 251, causes a sustained reduction of daily food intake in the rat. *Physiology and Behavior* **82** 863–869.
- Chard T 1987 An introduction to radioimmunoassay and related techniques. In *Laboratory Techniques in Biochemistry and Molecular Biology*, pp 1–74. Eds RH Burdon & H van Kinooenberg. Amsterdam: Elsevier Science.
- Devane WA, Dysarz FA, Johnson MR, Melvin LS & Howlett AC 1988 Determination and characterization of a cannabinoid receptor in rat-brain. *Molecular Pharmacology* **34** 605–613.
- Fernandez-Ruiz J, Munoz RM, Romero J, Villanua MA, Makriyannis A & Ramos JA 1997 Time course of the effects of different cannabimimetics on prolactin and gonadotrophin secretion: evidence for the presence of CB1 receptors in hypothalamic structures and their involvement in the effects of cannabimimetics. *Biochemical Pharmacology* **53** 1919–1927.
- Fernandez-Solari J, Scorticati C, Mohn C, De Laurentis A, Billi S, Franchi A, McCann SM & Rettori V 2004 Alcohol inhibits luteinizing hormone-releasing hormone release by activating the endocannabinoid system. *PNAS* **101** 3264–3268.
- Foster-Schubert KE & Cummings DE 2006 Emerging therapeutic strategies for obesity. *Endocrine Reviews* **27** 779–793.
- Gatley SJ, Gifford AN, Volkow ND, Lan RX & Makriyannis A 1996 I-123-labeled AM251: a radioiodinated ligand which binds *in vivo* to mouse brain cannabinoid CB1 receptors. *European Journal of Pharmacology* **307** 331–338.
- Hillard CJ, Farber NE, Hagen TC & Bloom AS 1984 The effects of delta-9-tetrahydrocannabinol on serum thyrotropin levels in the rat. *Pharmacology, Biochemistry, and Behavior* **20** 547–550.
- Klein TW 2005 Cannabinoid-based drugs as anti-inflammatory therapeutics. *Nature Reviews. Immunology* **5** 400–411.
- Lomax P, Kokka N & George R 1970 Thyroid activity following intracerebral injection of morphine in rat. *Neuroendocrinology* **6** 146–152.
- Di Marzo V & Fontana A 1995 Anandamide, an endogenous cannabinomimetic eicosanoid: killing two birds with one stone. *Prostaglandins, Leukotrienes, and Essential Fatty Acids* **53** 1–11.
- Di Marzo V, Bifulco M & De Petrocellis L 2004 The endocannabinoid system and its therapeutic exploitation. *Nature Reviews. Drug Discovery* **3** 771–784.
- Matias I & Di Marzo V 2007 Endocannabinoids and the control of energy balance. *Trends in Endocrinology and Metabolism* **18** 27–37.
- Matsuda LA, Lolait SJ, Brownstein MJ, Young AC & Bonner TI 1990 Structure of a cannabinoid receptor and functional expression of the cloned cDNA. *Nature* **346** 561–564.
- McLaughlin PJ, Winston KM, Limebeer CL, Parker LA, Makriyannis A & Salamone JD 2005 The cannabinoid CB1 antagonist AM251 produces food avoidance and behaviors associated with nausea but does not impair feeding efficiency in rats. *Psychopharmacology* **180** 286–293.
- de Miguel R, Romero J, Munoz R, Garcia-Gil L, Gonzalez S, Villanua MA, Makriyannis A, Ramos JA & Fernandez-Ruiz JJ 1998 Effects of cannabinoids on prolactin and gonadotrophin secretion: involvement of changes in hypothalamic gamma-aminobutyric acid (GABA) inputs. *Biochemical Pharmacology* **56** 1331–1338.

- Munro S, Thomas KL & Abushaar M 1993 Molecular characterization of a peripheral receptor for cannabinoids. *Nature* **365** 61–65.
- Oliveira KJ, Paula GS, Costa-e-Sousa RH, Souza LL, Moraes DC, Curty FH & Pazos-Moura CC 2006 Peptide YY (PYY)_{3–36} modulates thyrotropin secretion in rats. *Journal of Endocrinology* **191** 459–463.
- Ortiga-Carvalho TM, Polak J, McCann S & Pazos-Moura CC 1996 Effect of thyroid hormones on pituitary neuromedin B and possible interaction between thyroid hormones and neuromedin B on thyrotropin secretion. *Regulatory Peptides* **67** 47–53.
- Ortiga-Carvalho TM, Oliveira KJ, Soares BA & Pazos-Moura CC 2002 The role of leptin in the regulation of TSH secretion in the fed state: *in vivo* and *in vitro* studies. *Journal of Endocrinology* **174** 121–125.
- Pagotto U, Marsicano G, Fezza F, Theodoropoulou M, Grubler Y, Stalla J, Arzberger T, Milone A, Losa M, Di Marzo V *et al.* 2001 Normal human pituitary gland and pituitary adenomas express cannabinoid receptor type 1 and synthesize endogenous cannabinoids: first evidence for a direct role of cannabinoids on hormone modulation at the human pituitary level. *Journal of Clinical Endocrinology and Metabolism* **86** 2687–2696.
- Pazos-Moura CC, Moura EG, Rettori V, Polak J & McCann S 1996 Role of neuromedin B in the *in vitro* thyrotropin release in response to thyrotropin-releasing hormone (TRH) from anterior pituitaries of eu-, hypo- and hyperthyroid rats. *Proceedings of the Society for Experimental Biology and Medicine* **211** 353–358.
- Pertwee RG 2005 Inverse agonism and neutral antagonism at cannabinoid CB1 receptors. *Life Sciences* **76** 1307–1324.
- Porcella A, Marchese G, Casu MA, Rocchitta A, Lai ML, Gessa GL & Pani L 2002 Evidence for functional CB1 cannabinoid receptor expressed in the rat thyroid. *European Journal of Endocrinology* **147** 255–261.
- Rettori V, Milenkovic L, Fahim AM, Polak J, Bloom SR & McCann SM 1989 Role of neuromedin B in the control of the release of thyrotropin in the rat. *PNAS* **86** 4789–4792.
- Rettori V, Pazos-Moura CC, Moura EG, Polak J & McCann SM 1992 Physiological role of neuromedin B in control release of thyrotropin in hypothyroid and hyperthyroid rats. *PNAS* **89** 3035–3039.
- Scorticati C, Mohn C, De Laurentis A, Vissio P, Solari JF, Seilicovich A, McCann SM & Rettori V 2003 The effect of anandamide on prolactin secretion is modulated by estrogen. *PNAS* **100** 2134–2139.
- Scorticati C, Fernandez-Solari J, De Laurentis A, Mohn C, Prestifilippo JP, Lasaga M, Seilicovich A, Billi S, Franchi A, McCann SM *et al.* 2004 The inhibitory effect of anandamide on luteinizing hormone-releasing hormone secretion is reversed by estrogen. *PNAS* **101** 11891–11896.
- Tsou K, Brown S, Sanudo-Pena MC, Mackie K & Walker JM 1998 Immunohistochemical distribution of cannabinoid CB1 receptors in the rat central nervous system. *Neuroscience* **83** 393–411.
- Weidenfeld J, Feldman S & Mechoulam R 1994 Effect of the brain constituent anandamide, a cannabinoid receptor agonist, on the hypothalamo-pituitary-adrenal axis in the rat. *Neuroendocrinology* **59** 110–112.
- Weintraub BD, Kazlauskaitė R, Grossmann M & Szkudlinski MW 2001 Thyroid-stimulating hormone and regulation of the thyroid axis. In *Endocrinology*, edn 4, pp 1345–1360. Eds L DeGroot & JL Jameson. Philadelphia: Saunders Company.
- Wenger T, Toth BE & Martin BR 1995 Effects of anandamide (endogenous cannabinoid) on anterior-pituitary hormone-secretion in adult ovariectomized rats. *Life Sciences* **56** 2057–2063.
- Wenger T, Jamali KA, Juaneda C, Léonardelli J & Tramu G 1997 Arachidonyl ethanolamide (anandamide) activates the parvocellular part of hypothalamic paraventricular nucleus. *Biochemical and Biophysical Research Communications* **237** 724–728.
- Wenger T, Fernandez-Ruiz JJ & Ramos JA 1999 Immunocytochemical demonstration of CB1 cannabinoid receptors in the anterior lobe of the pituitary gland. *Journal of Neuroendocrinology* **11** 873–878.
- Xi ZX, Gilbert JG, Peng XQ, Pak AC, Li X & Gardner EL 2006 Cannabinoid CB receptor antagonist AM251 inhibits cocaine-primed relapse in rats: role of glutamate in the nucleus accumbens. *Journal of Neuroscience* **26** 8531–8536.
- Zar J 1996 Data transformations. In *Biostatistical Analysis*, 3rd edn, pp 277–279. London: Prentice Hall International.

Received in final form 20 August 2008

Accepted 27 August 2008

Made available online as an Accepted Preprint

28 August 2008